

# Management of Twin Pregnancies (Part 1)

## PRINCIPAL AUTHORS

Jon Barrett (Editor and Chair), MD, FRCSC  
Alan Bocking (Co-chair), MD, FRCSC

### Subgroup A: MULTIPLE BIRTH 0-20 WEEKS. RECOMMENDATIONS FOR EARLY PREGNANCY ULTRASOUND AND GENETIC COUNSELLING

Hani Akoury (Coordinator), MD, FRCSC, Toronto, ON  
Jo-Anne Johnson (Coordinator), MD, FRCSC, Toronto, ON  
Nan Okun, MD, FRCSC, Edmonton, AB  
Anne Pastuzak, PhD, Toronto, ON  
Gareth Seaward, MD, FRCSC, Toronto, ON  
J. Wilkinson, MD, FRCSC, Toronto, ON  
Doug Wilson, MD, FRCSC, Vancouver, BC

### Subgroup B: PRETERM LABOUR PREVENTION

Tony Armson (Coordinator), MD, FRCSC, Halifax, NS  
André Bastide, MD, FRCSC, Quebec City, QC  
Joan Crane, MD, FRCSC, St. John's, NF  
Robert Goldenberg, MD, Birmingham, AB, USA  
Mary Hannah, MD, FRCSC, Toronto, ON  
John Kingdom, MD, FRCSC, Toronto, ON  
Steve Lye, PhD, Toronto, ON  
Renato Natale, MD, FRCSC, London, ON  
David Rouselle, MD, FRCSC, Newmarket, ON  
Rory Windrim, MD, FRCSC, Toronto, ON

### Subgroup C: FETAL GROWTH IN MULTIPLES

Line Leduc (Coordinator), MD, FRCSC, Montreal, QC  
John Dube, MD, FRCSC, Halifax, NS  
Dan Farine, MD, FRCSC, Toronto, ON  
Kathy Fong, MD, FRCSC, Toronto, ON  
Louis Keith, MD, Chicago, USA  
Ian Lange, MD, FRCSC, Calgary, AB  
David Rosman, MD, FRCSC, Toronto, ON  
David Young, MD, FRCSC, Halifax, NS

## Subgroup D:

### LABOUR MANAGEMENT AND THE DELIVERY OF THE SECOND TWIN— THE EVIDENCE AND RECOMMENDATION

Elysse Levinsky (Coordinator), MD, FRCSC, Toronto, ON  
Peter Von Daedelzen (Coordinator), MD, MRCOG, Toronto, ON  
Kofi Amankwah, MD, FRCSC, Toronto, ON  
Greg Davies, MD, FRCSC, London, ON  
Michael Helewa, MD, FRCSC, Winnipeg, MB  
Jay McGilivray, Toronto, ON  
Ken Milne, MD, FRCSC, Ottawa, ON  
Knox Ritchie, MD, FRCSC, Toronto, ON

## Subgroup E:

### SPECIAL TWIN CIRCUMSTANCES—GUIDELINES FOR DIAGNOSIS AND MANAGEMENT (TWIN TWIN TRANSFUSION, MONOAMNIOTIC PREGNANCIES)

Karen Fung Kee Fung (Coordinator), Ottawa, ON  
Karen Ash, MD, FRCSC, Ottawa, ON  
Nick Fisk, MB, FRCOG, Toronto, ON  
André Gagnon, MD, FRCSC, Vancouver, BC  
Rob Gratton, MD, FRCSC, London, ON  
Geoff Machin, MD, PhD, Oakland, CA, USA  
Karel O'Brien, MD, FRCSC, Toronto, ON  
Greg Ryan, MD, FRCSC, Toronto, ON  
John Smith, MD, FRCSC, Hamilton, ON

## Subgroup F:

### THE IMPACT OF MULTIPLES—THE INCIDENCE, PERINATAL MORTALITY RATE (PNMR), BURDEN OF ILLNESS

Alan Bocking (Coordinator), MD, FRCSC, London, ON  
Elizabeth Asztalos, MD, FRCSC, Toronto, ON  
Elizabeth Bryan, MD, FRCP, London, UK  
Cathy Cameron, London, ON  
Catherine Harrison, Ottawa, ON  
Donna Launslager, Windsor, ON  
Catherine McCourt, MD, Ottawa, ON  
Patricia Niday, EdD, Ottawa, ON  
Jeff Nisker, MD, FRCSC, London, ON  
Paula Stewart, MD, Ottawa, ON  
Donna Wilson, RN, MN, Toronto, ON

These guidelines reflect emerging clinical and scientific advances as of the date issued and are subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Local institutions can dictate amendments to these opinions. They should be well documented if modified at the local level. None of the contents may be reproduced in any form without prior written permission of SOGC.

## INTRODUCTION

The incidence of twin gestations across Canada has increased by 15 percent over the years 1993-1997. This increase is the major contributing factor to the rising incidence of preterm birth in Canada.<sup>1</sup> Many aspects of the obstetric management of the twin pregnancy cannot be extrapolated from that of a singleton pregnancy. For example, the age related karyotypic risk is different for a twin compared to a singleton pregnancy;<sup>2</sup> during pregnancy the clinical assessment of the growth of each fetus is difficult without ultrasound; and finally, the delivery of the second twin demands special attention. Therefore in December 1997, the SOGC and the Universities of Toronto and Western Ontario\* coordinated a National Consensus Conference in order to define, based upon the best evidence available, the standard care of twin and higher order multiples gestation.

## PROCESS

Five priority areas were identified and questions relating to those areas developed in order to allow participants to concentrate on important guidelines for practice. The impact of the increase in multiple births was also addressed. The participants were divided into groups to address each of these topics. Each group had in charge a leader or leaders whose task it was to review the literature and provide evidence based answers and forward the group's consensus on particular issues. The quality of evidence was evaluated and recommendations were

made according to guidelines for assessing medical literature published by Health Canada (Table I).<sup>3</sup> As an aid to the reader, alongside each recommendation is the Level of Evidence and Category of Strength for that recommendation. Although many areas of practice have not been well studied, clinical practice dictates that some recommendations of practice are made to aid professionals in their day to day clinical activity. It is important to realize that these particular recommendations are made not on "best evidence" but on "best opinion" and should be flagged as areas for future research. The full deliberations of each group will be available on the SOGC "Multiples" Website and in a more complete publication which will be available in the future.

It was attempted to keep medical representation at the meeting wide, with input sought from all aspects of health care including midwifery, nursing, counselling, social work, as well as community groups. We are totally indebted to the group facilitators who reviewed and presented vast amounts of literature and who have allowed us to present this consensus document.

### A. THE INCIDENCE AND IMPACT OF TWIN PREGNANCIES

Multiple births have increased from 1.9 percent to 2.1 percent of all live births between 1981 through 1983 and 1992 through 1994. The contribution of these multiple pregnancies to preterm birth rates in Canada has resulted in a 25 percent increase in the proportion of preterm births resulting from multiple gestations.<sup>1</sup> The impact of this data on both our health care system and on Canadian society, along with the

TABLE 1 <sup>3</sup> QUALITY OF EVIDENCE ASSESSMENT	CLASSIFICATION OF RECOMMENDATIONS
<p>The quality of evidence reported in these guidelines has been described using the Evaluation of Evidence criteria outlined in the Report of the Canadian Task Force on the Periodic Health Exam.<sup>3</sup></p> <p>I: Evidence obtained from at least one properly randomized controlled trial.</p> <p>II-1: Evidence from well-designed controlled trials without randomization.</p> <p>II-2: Evidence from well-designed cohort (prospective or retrospective) or case-control studies, preferably from more than one centre or research group.</p> <p>II-3: Evidence obtained from comparisons between times or places with or without the intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940's) could also be included in this category.</p> <p>III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees.</p>	<p>Recommendations included in these guidelines have been adapted from the ranking method described in the Classification of Recommendations found in the Report of the Canadian Task Force on the Periodic Health Exam.<sup>3</sup></p> <p>A. There is good evidence to support the recommendation that the condition be specifically considered in a periodic health examination.</p> <p>B. There is fair evidence to support the recommendation that the condition be specifically considered in a periodic health examination.</p> <p>C. There is poor evidence regarding the inclusion or exclusion of the condition in a periodic health examination, but recommendations may be made on other grounds.</p> <p>D. There is fair evidence to support the recommendation that the condition not be considered in a periodic health examination.</p> <p>E. There is good evidence to support the recommendation that the condition be excluded from consideration in a periodic health examination.</p>

\* With support of the Medical Research Council of Canada, Adeza, Ferring, Hewlett Packard, Mantria Health Care, Serono, and Upjohn

recommendations of the subgroup that addressed these issues, will be published in the August issue of *Journal SOGC*.

## B. RECOMMENDATIONS FOR EARLY PREGNANCY ULTRASOUND AND GENETIC COUNSELLING.

### 1) The first trimester ultrasound in twin pregnancy?

Chorionicity, one of the most important determinants of pregnancy outcome in twin gestation, is best determined in the first trimester. Nuchal translucency can also be measured.

#### CONSENSUS STATEMENT #1

##### **When a multiple gestation has been diagnosed:**

- Every effort should be made to determine chorionicity at the time of diagnosis. (II-3 C)
- The optimal time to determine chorionicity is 10-14 weeks. (II-3 C)
- While these recommendations apply to diagnosis of twin pregnancy with regard to prenatal diagnosis and counselling, there have been no studies relating the establishment of prenatal chorionicity to pregnancy outcome.

### 2) At what age should genetic testing be offered to a mother with dichorionic and monochorionic twins?

Invasive diagnostic testing may be offered in twins on the basis of late maternal age. When counselling women about their risk of chromosome abnormalities, the chorionicity should be taken into consideration. In monochorionic (MC) twins, the age-related risk for the fetuses is the same (all are monozygotic) and is equivalent to the risk in a singleton pregnancy. In dichorionic (DC) twins, the risk is essentially double the age-related risk (about 2/3 will be dizygotic). Although accurate determination of chorionicity by ultrasound is possible, it will not always be feasible. Hence, the group felt it was more appropriate at this time to follow the guidelines by Rodis *et al.* (1990) which state: "The chance of a 32 year old woman who carries twins of unknown zygosity having at least one child with Down syndrome is equivalent to the risks of a 35 year old with a singleton pregnancy."<sup>2</sup>

The risk of amniocentesis in twin gestation is uncertain (see below) and issues such as discordant anomalies that may arise need to be considered. Such counselling is complicated and is best carried in a specialized genetic centre, or centre specializing in the management of multiple gestation.

#### CONSENSUS STATEMENT #2

**All women carrying twin pregnancies should be referred for counselling to a centre for the consideration of invasive testing at age 32. The counselling must be individualized and the final decision must be taken by the parents since the risk of amniocentesis is uncertain in twin gestation. (II-3 C)**

### 3) What are the methods available for genetic screening in twin pregnancies and how effective are they in twin pregnancies in detecting fetal aneuploidy?

#### CONSENSUS STATEMENT #3

- Biochemical screening for aneuploidy is not recommended in twins.
- Maternal serum alpha fetoprotein (MS-AFP) is useful for detection of open neural tube and other birth defects. (II-3 C)
- Evidence is promising that nuchal translucency (NT) screening is useful for identifying twin pregnancies at high risk of aneuploidy. This requires further prospective investigation. (II-3 C)

### 4) What are the risks/benefits of invasive genetic testing in twin pregnancy?

#### CONSENSUS STATEMENT #4

- The fetal loss rates with invasive testing (amniocentesis and chorionic villus sampling (CVS)) in twins are unclear. (II-3 C)
- Development of a protocol for standardization of technique (as determined by expert opinion) is recommended.
- Invasive testing should be offered to twins according to the usual standard of care.

#### CONSENSUS STATEMENT #5

- In twins discordant for abnormality, the option of selective reduction should be offered. The procedure should be performed in a tertiary level center. Transportation and out-of-province costs should be covered.

## C. PRETERM BIRTH PREVENTION IN TWIN GESTATION.

The use of tocolytics and corticosteroids in the treatment of established preterm labour was not addressed at this meeting. Readers are directed to *The Canadian Consensus on the Use of Tocolytics for Pre-Term Labour*.<sup>4</sup>

### 1) Is there any evidence that bedrest, cervical suture or tocolysis, or any other intervention prevents preterm labour and delivery in twin pregnancies?

#### HOSPITAL BEDREST

Randomized controlled trials and a meta-analysis of hospital bedrest in twin pregnancies have shown no reduction in preterm birth or perinatal death.<sup>5-12</sup> In uncomplicated twin pregnancies, hospital rest may result in increased risk of very preterm birth and maternal psychosocial stress. In women with twin pregnancy at high risk for preterm birth because of premature cervical change prior to labour, there is no evidence that hospital bedrest will reduce the rate of preterm birth.

---

CONSENSUS STATEMENT #6

**Routine hospitalization for bedrest in multiple gestation is *not* recommended. (I E)**

ACTIVITY RESTRICTION/WORK LEAVE

Restriction of activity level and the recommendation to stop work is commonly prescribed for women with twin pregnancies as a preterm birth prevention strategy. This prophylactic intervention has only been studied in a few observational trials with historical or geographic controls with conflicting results.<sup>13,14</sup>

CONSENSUS STATEMENT #7

**There is insufficient evidence to support prophylactic activity restriction or work leave in multiple gestation. (III C)**

CERVICAL CERCLAGE

Prophylactic cervical cerclage has not been shown to be effective in preventing preterm birth in twin pregnancy in observational or controlled trials.<sup>15</sup>

CONSENSUS STATEMENT #8

**There is moderate evidence against routine prophylactic cervical cerclage in multiple gestation. However, cerclage may be indicated for the treatment of incompetent cervix or other specific circumstances. (I;II-2 D)**

PROPHYLACTIC TOCOLYTIC THERAPY

Most randomized controlled trials have failed to show any benefit of prophylactic oral or intravenous tocolytic therapy in multiple gestation.<sup>16-20</sup>

CONSENSUS STATEMENT #9

**There is moderate evidence against prophylactic tocolysis in the management of multiple gestation, but it may be indicated on other grounds. (I;II-2 D)**

SPECIALIZED TWIN CLINICS/PREVENTION PROGRAMMES

Multi-intervention preterm birth prevention programmes for twin pregnancies have been evaluated in several observational studies with contemporary and/or historical controls.<sup>20-26</sup> All studies reviewed suggest reduction in preterm birth rate, decreased perinatal mortality, and overall improvement in perinatal outcome.

CONSENSUS STATEMENT #10

**The evidence to support specialized clinics is of insufficient quality to recommend that they be part of routine clinical practice. Further randomized controlled studies are needed to validate the improved outcomes that have been**

**demonstrated in cohort studies. (II-2;III C)**

**2) Is there a role for routine clinical cervical assessment in multiple gestation?**

Despite the lack of precision, clinical cervical assessment appears to be safe and may be effective in monitoring twin gestations, if transvaginal ultrasound is not available or determined to be too expensive.<sup>27</sup> However, compared to transvaginal sonography, digital examination is more subjective and less reproducible.<sup>28-30</sup>

CONSENSUS STATEMENT #11

**There is good evidence that premature cervical change by digital examination predicts preterm birth in twins. (II-2 A) Since there are no well designed intervention trials available, the role of sonographic clinical cervical assessment in the prenatal period has not been determined. (C)**

**3) Is there a role for sonographic cervical assessment in multiple gestation?**

Transvaginal sonographic cervical assessment provides insight into the cervical status, as well as the likelihood of preterm birth in twin pregnancies. There appears to be good correlation between cervical length and the risk of preterm birth.<sup>31-35</sup>

CONSENSUS STATEMENT #12

**There is good evidence that transvaginal sonographic measurement of cervical length predicts preterm birth in twins. (II-1 A) While the predictive ability of cervical length measurement is established, there are no intervention studies that have evaluated cervical length measurement in the prevention of preterm birth, and therefore the role of sonographic clinical cervical assessment in the prenatal period has not been determined. (C)**

**4) Is home uterine activity monitoring useful in predicting preterm birth in twin pregnancy?**

Although home uterine activity monitoring may be helpful in identifying women at increased risk of preterm labour before advanced cervical dilation occurs, this information has not resulted in reduction in the incidence of preterm labour, advanced cervical dilation at presentation or preterm birth in well-controlled randomized controlled trials.<sup>36-40</sup>

CONSENSUS STATEMENT #13

**There is moderate evidence against home uterine activity monitoring in multiple gestation. (I D)**

**5) Does the measurement of fetal fibronectin predict preterm birth in twin pregnancy?**

Data from prospective longitudinal studies suggests that a positive fetal fibronectin test has a very high negative predictive value for the prediction of preterm birth in asymptomatic

patients. The positive predictive value for preterm labour and delivery before 37 weeks is 60 percent for patients in preterm labour, 45 percent in asymptomatic high-risk women, and 30 percent in asymptomatic low-risk women.<sup>41</sup>

#### CONSENSUS STATEMENT #14

**There is good evidence that the presence of cervicovaginal fetal fibronectin in twins predicts preterm birth. Without well designed intervention trials available, there is no basis for incorporating fetal fibronectin screening into routine prenatal management of multiple gestation. (C)**

#### SUMMARY

No prenatal method has been shown to prevent preterm birth labour and birth in twin pregnancies. Cervical length measurement and presence of fetal fibronectin in cervico-vaginal fluids need to be evaluated to investigate whether the ability to predict whether selected interventions, applied to a subgroup of twins with a high risk of preterm birth, will result in a reduction in the rate of preterm birth.

#### D. ULTRASOUND IN TWIN GESTATIONS<sup>42-69</sup>

##### 1) When should ultrasounds be performed in twin pregnancies and why?

Without ultrasound, up to 40 percent of twin gestations will not be recognized until 26 weeks gestation on average, and up to 20 percent will remain unrecognized until term. In the first or second trimester ultrasound will usually (>95%) determine chorionicity. The detection of fetal anomaly, the incidence of which is three times higher with twin pregnancy, is initially best assessed between 16 and 20 weeks. In the second and third trimesters, fetal growth will be reliably assessed by serial ultrasound.

#### CONSENSUS STATEMENT #15

**There is good evidence that the diagnosis of twin gestation is improved by the routine use of ultrasound. There is consensus that serial ultrasonographic evaluation every three to four weeks is indicated in twin gestations. (I B)**

##### 2) Is fetal growth the same in twins as in singletons and what ultrasonic growth curves should be used to plot fetal weights?

Fetal growth in twin gestation parallels that of singletons until approximately 32-35 weeks. Thereafter the rate of fetal growth is measurably slightly less, although the clinical significance of this is undetermined. The patterns of twin fetal growth vary by race and gender, with African-American mothers having lower median body weight (BW) values. Male twins have heavier median BW for gestational age than female twins at every gestational age.

#### CONSENSUS STATEMENT #16

**Fetal growth differs slightly in twin gestations and twin specific charts may be used to define the normal growth rate. Precision may also be obtained by using sex and race specific charts. In clinical practice, however, these differences are small and singleton growth curves may be used. Patterns of fetal growth are more important than absolute measurements. Both must be interpreted in the light of the clinical history, together with all the genetic and environmental factors that may affect fetal growth. (III B)**

##### 3) What level of growth discrepancy between twins is cause for concern?

The importance of accurate and timely identification of discordant growth lies primarily in its relationship to the complication of twin twin transfusion and to intra-uterine growth restriction (IUGR) of the smaller twin. True discordance is an indicator for an increased risk of IUGR, morbidity, and mortality for the smaller twin. A risk for aneuploidy, anomaly or viral syndrome affecting only one fetus must also be considered when discordant growth is identified.

#### CONSENSUS STATEMENT #17

**The diagnosis of discordance has been based on the following:**

- an abdominal circumference (AC) difference of 20 mm (sensitivity of 80%, specificity 85%, positive predictive value (PPV) 62%)
- estimated fetal weight (EFW) based on bi-parietal diameter (BPD) and AC or AC and femur length (FL) > 20 percent (sensitivity 25-55%) (II-2 B)

#### E. LABOUR MANAGEMENT AND THE DELIVERY IN TWIN GESTATION

##### 1) What are the indications for elective Caesarean in twin pregnancies (> 2,500g)?

#### CONSENSUS STATEMENT #18

**The indications for elective Caesarean section in twin gestations are:**

- a) **Monoamniotic twins** because the risk of entrapment is too great to permit elective vaginal delivery;
- b) **Conjoined twins** other than at gestations remote from term;
- c) Indications as for singleton pregnancies. (III C)

##### 2) What are the attendance at labour and delivery guidelines for physicians managing a twin gestation?

#### CONSENSUS STATEMENT #19

**The following points highlight the factors most crucial to**

---

### **the successful care of a woman during labour and delivery of a twin pregnancy.**

- a) Timely attendance by a physician competent to manage a twin birth.
- b) The presence of additional antenatal risk factors should be reviewed at the onset of labour. Intrapartum risk factors should be assessed on an ongoing basis and changes attended to appropriately.
- c) When participating in a call system, the replacing physician should be of similar competence and informed of all facts pertaining to a case when care is transferred.
- d) The diagnosis of twins is usually antenatal. Therefore, arrangements for delivery and/or transfer should be set in place. This may include antenatal consultation with a high risk centre.
- e) The assessment of lie and presentation of each fetus on admission in labour, preferably by ultrasound.
- f) Intravenous access should be secured, and blood sent for group and antibody screen.
- g) Anaesthetic personnel should be informed as soon as possible of a planned twin birth. Epidural anaesthesia is advantageous.
- h) Oxytocin augmentation may be used before the delivery of the first twin and/or between twins for hypotonic contractions.
- i) For either twin, the indication(s) for any intervention should be convincing, compelling, and documented at the time of the event(s). However, for the cephalic second twin, vaginal delivery should be expedited should fetal distress occur. There is little evidence to suggest the best operative method of delivering the second twin who remains in the vertex position, should the need arise. A vacuum or forceps procedure, preferably during which the prerequisites of any operative delivery are fulfilled, may be considered. However, the vacuum may be employed at a station perhaps slightly higher than would be attempted with a singleton fetus. Other options are conversion to breech and delivery by breech extraction if the Vx was not engaged or Caesarean if the operator felt that no other method could be safely accomplished.
- j) Documentation of all aspects of labour and delivery should be clear, contemporaneous, and consistent among all involved health care providers.
- k) Progress of labour should emerge clearly from the documentation.
- l) Continuous electronic fetal heart rate monitoring of both twin A and B should ensure that both twins are being monitored individually. The presence of an ultrasound machine in the delivery room may be advantageous.
- m) For attempted delivery by mid-forceps, vaginal breech delivery, and multiple pregnancies, Caesarean section should be available immediately. Immediate availability means the presence in the hospital of anaesthetic, obstetrical, neonatal, and nursing staff trained in Caesarean delivery. A note should be dictated describing all operative deliveries and complicated labour and delivery events. The time difference

between the delivery of each baby should be noted.

- n) Cord blood samples should be drawn at the time of delivery.
- o) The third stage of labour should be managed actively, with oxytocin being administered with the delivery of the second twin.
- p) Placentas should undergo gross and microscopic pathological examination.
- q) We suggest that twin deliveries be planned in Level II and Level III hospitals. (II C)

### **3) What is the best method of delivering the non vertex second twin?**

#### **CONSENSUS STATEMENT #20**

- a) Delivery of cephalic twin A/non-cephalic twin B: Estimated weight 1,500-4,000 g. Vaginal delivery is indicated as long as the obstetrician is comfortable with and skilled in vaginal breech delivery.<sup>70-83</sup> (II-2 B)
- b) Delivery of cephalic twin A/non-cephalic twin B: Estimated weight 500-1,500 g. In this weight range the group acknowledged that there is no consistent evidence to support either Caesarean section or the vaginal route for delivery.<sup>70-78</sup> (III C)

### **4) Non-cephalic twin B: Breech extraction with or without internal podalic version OR external cephalic version?**

#### **CONSENSUS STATEMENT #21**

**Breech extraction with or without internal podalic version is associated with a lower Caesarean section rate and similar neonatal and maternal outcomes compared with external cephalic version in the twin pairs whose estimated fetal weights are greater than 1,500 g.<sup>79-84</sup> (II-2 B)**

### **5) What is the optimal delivery interval between twins in a vaginal twin delivery?**

#### **CONSENSUS STATEMENT #22**

**It is reasonable to either:**

- a) expedite delivery by oxytocin infusion, stabilizing amniotomy and, if indicated, operative vaginal delivery; or
- b) permit a longer delay between deliveries using an oxytocin infusion and continuous electronic fetal heart rate monitoring. For the non-cephalic second twin, if breech extraction with or without podalic version is considered, then this should be done without delay.<sup>85-88</sup> (II-2 B)

### **F. Special Twin Circumstances, Guidelines For Diagnosis And Management: Twin Twin Transfusion (TTTS), Monoamniotic Pregnancies.**

#### **1) What are the diagnostic markers of twin twin transfusion?**

CONSENSUS STATEMENT #23

**a) Evidence for diagnosis:<sup>89-111</sup>**

- i) single monochorionic placenta,
- ii) polyhydramnios/oligohydramnios sequence, and
- iii) same sex fetuses.

**This does not imply that all pregnancies characterized by these features are affected by TTTS—further investigations may be required—but these features should prompt referral to a tertiary care unit. (II-2 B)**

**b) Evidence of decompensation in fetal health. These include:**

- i) chronically distended bladder of recipient twin;
- ii) growth discordance (>25%); and
- iii) evidence of cardiac dysfunction eg., non-immune hydrops.

These should prompt urgent referral to a tertiary care perinatal centre. (II-2 B)

**2) How should TTTS pregnancies be managed antenatally and when should they be delivered?**

CONSENSUS STATEMENT #24

**In consideration of the high perinatal mortality rate accompanying a diagnosis of TTTS, all monochorionic twin pregnancies should be seen in consultation by a maternal-fetal medicine specialist for counselling and monitoring of fetal condition. Urgent consultation should be sought for the criteria listed in Section 23(b) above. (II-3 B)**

**a) Monitoring (uncomplicated monochorionic twin pregnancy):**

Following diagnosis of uncomplicated monochorionic twin pregnancy and detailed second trimester anomaly screen, serial ultrasound surveillance of fetal health performed every two weeks, looking for evidence of suboptimal growth and/or emergence of features suggestive of TTTS. (III C)

**b) Established TTTS:**

The frequency of ultrasound surveillance of fetal health in severe cases of TTTS will depend on the severity and intervention strategy. (III C)

**Therapeutic options: These include:**

- i) no intervention (survival 0—30%),
- ii) amnioreduction 64 percent (survival 64% overall, 74% of at least one twin),
- iii) laser photocoagulation (55% overall survival—73% of at least one twin),
- iv) amniotic septostomy, 83 percent survival (12 cases only). Termination of pregnancy, either selective or total may be considered.<sup>112-137</sup>

**3) What are the diagnostic features and risks of monoamniotic twin gestation?**

CONSENSUS STATEMENT #25

**Monochorionic, monoamniotic placentation is found in approximately 1 percent of all twin gestations. High mortality rates (up to 50%) have been attributed to cord entanglement, knots and twists, congenital anomalies, and prematurity. Cord entanglement is present at birth in virtually 100 percent of monoamniotic twins and is a key diagnostic feature of monoamnioticity on sonography. Other sonographic diagnostic features of a monoamniotic twin pregnancy include:**

- a) absence of a dividing amniotic membrane,
- b) presence of a single placenta,
- c) both fetuses of the same gender,
- d) adequate amniotic fluid surrounding each fetus, and
- e) both fetuses moving freely within the uterine cavity.<sup>138-156</sup>

**4) How should they be managed antenatally and when should they be delivered?**

CONSENSUS STATEMENT #26

**These cases should all be referred to a regional perinatal centre. A reasonable management plan includes:**

**Antenatal:**

- a) frequent (weekly or more) non stress test commencing at 24 weeks, (II-3)
- b) judicious use of antenatal corticosteroids. (III C)

**Delivery:**

- a) By 32 -33 weeks, although some evidence that later delivery may be appropriate, (II-3,2 C)
- b) Casearean delivery.<sup>145,150,153, 156</sup> (II B)

J Soc Obstet Gynaecol Can 2000;22(7):519-29

**REFERENCES**

1. Josephs KS, Kramer MS, Wu Wen S, Alexander D. Determinants of pre term birth in Canada from 1981 through 1983 and 1992 through 1994. N Engl J Med 1998;339:1434-9.
2. Rodis JF, Egan, Craffey A., Ciarleglio, L. Greenstien, RH., Scorza, WE. Calculated Risk of Chromosomal Abnormalities in Twin Gestations. Obstet Gynecol 1990 76:1037-41.
3. Woolf SH, Battista RN, Angerson GM, Logan AG, EEL W. Canadian Task Force on the Periodic Health Exam. Ottawa: Canada Communication Group, 1994:xxxvii.
4. Hannah M (Chair). The Canadian Consensus on the Use of Tocolytics for Pre- Term Labour. J Soc Obstet Gynaecol Can 1996;17,11:1089.
5. Crowther CA. Hospitalization for bed rest in multiple pregnancy (Cochrane Review). In: The Cochrane Library, Issue 3, 1998. Oxford: Update Software.

6. Hartikainen-Sorri AL, Jouppila P. Is routine hospitalization needed in antenatal care of twin pregnancy? *J Perinat Med* 1984;12:31-4.
7. Saunders MC, Dick JS, Brown IM, McPherson K, Chalmers I. The effects of hospital admission for bed rest on the duration of twin pregnancy: a randomized trial. *Lancet* 1985;2:793-5.
8. MacLennan AH, Green RC, O'Shea R, Brookes C, Morris D. Routine hospital admission in twin pregnancy between 26 and 30 weeks' gestation. *Lancet* 1990;335:267-9.
9. Crowther CA, Neilson JP, Verkuyl DAA, Bannerman C, Ashurst HM. Preterm labour in twin pregnancies: can it be prevented by hospital admission? *Br J Obstet Gynaecol* 1989;96:850-3.
10. Crowther CA, Verkuyl DAA, Neilson JP, Bannerman C, Ashurst HM. The effects of hospitalization for rest on fetal growth, neonatal morbidity and length of gestation in twin pregnancy. *Br J Obstet Gynaecol* 1990;97:872-7.
11. Younis JS, Sadovsky E, Eldar-Geva T, Mildwidsky A, Zeevi D, Zajicek G. Twin gestations and prophylactic hospitalization. *Int J Gynaecol Obstet* 1990;32:325-30.
12. Al-Najashi SS, Mulhim AA. Prolongation of pregnancy in multiple pregnancy. *Int J Gynaecol Obstet* 1996;54:131-5.
13. Papiernik E, Richard A, Tafforeau J, Keith L. Social groups and prevention of preterm births in a population of twin mothers. *J Perinat Med* 1996;24:669-76.
14. Rydhström H. Twin pregnancy and the effects of prophylactic leave of absence on pregnancy duration birth weight. *Acta Obstet Gynecol Scand* 1988;67:81-4.
15. Zakut H, Insler V, Serr DM. Elective cervical suture in preventing premature delivery in multiple pregnancies. *Israel J Med Sci* 1977;13:488.
16. Cetrulo CL, Freeman RK. Ritrodine HCL for the prevention of premature labor in twin pregnancies. *Acta Genet Med Gemellol Roma* 1976;321-4.
17. Marivate M, DeVilliers KQ, Fairbrother P. Effect of prophylactic outpatient administration of fenoterol on the time of onset of spontaneous labor and fetal growth rate in twin pregnancy. *Am J Obstet Gynecol* 1977;128:707.
18. O'Connor MC, Murphy H, Dalrymple IJ. Double blind trial of ritrodine and placebo in twin pregnancy. *Br J Obstet Gynecol* 1979;86:706-9.
19. Gummerus M, Halonen O. Prophylactic long-term oral tocolysis of multiple pregnancies. *Br J Obstet Gynaecol* 1987;94:249-51.
20. Ashworth MF, Spooner SF, Verkuyl DA, Waterman R, Ashurst HM. Failure to prevent preterm labour and delivery in twin pregnancy using prophylactic oral salbutamol [see comments]. *Br J Obstet Gynaecol* 1990;97:878-82.
21. Tafforeau J, Papiernik E, Richard A, Pons JC. Is prevention of preterm births in twin pregnancies possible? Analysis of the results of a prevention program in France (1989-1991). *Eur J Obstet Gynecol Reprod Biol* 1995;59:169-74.
22. Papiernik E, Mussy MA, Vial M, *et al.* A low rate of perinatal deaths for twin births. *Acta Genet med Gemellol* 1985;34:201-6.
23. D'Alton ME, Malozzi A, Dudley DK. The prevention of preterm delivery in twin pregnancy. Proceedings, Society of Perinatal Obstetricians, Eighth Annual meeting, Las Vegas, February, 1988.
24. Vergani P, Ghidini A, Bozzo G, *et al.* Prenatal management of twin gestation: Experiences with a new protocol. *J Reprod Med* 1991;36:667-71.
25. Plank K, Mikulaj V, Stencil J, Drobna H, Klesken P. Prevention and treatment of prematurity in twin gestation. *J Perinat Med* 1993;21:309-13.
26. Newman RB, Gill PJ, Katz M. Uterine activity during pregnancy in ambulatory patients: comparison of singleton and twin gestations. *Am J Obstet Gynecol* 1986;154:530.
27. Bivins HA Jr, Newman RB, Ellings JM, Hulse TC, Keenan A. Risks of antepartum cervical examination in multifetal gestations. *Am J Obstet Gynecol* 1993;169:22-5.
28. Newman RB, Godsey RK, Ellings JM, Campbell BA, Eller DP, Miller MC. Quantification of cervical change: relationship to preterm delivery in the multifetal gestation. *Am J Obstet Gynecol* 1991;165:264-9; discussion 269-71.
29. Gomez R, Galasso M, Romero R, *et al.* Ultrasonographic examination of the uterine cervix is better than cervical digital examination as a predictor of the likelihood of premature delivery in patients with preterm labor and intact membranes. *Am J Obstet Gynecol* 1994;171:956-64.
30. Neilson JP, Verkuyl DA, Crowther CA, Bannerman C. Preterm labor in twin pregnancies: prediction by cervical assessment. *Obstet Gynecol* 1988;72:719-23.
31. Kushnir O, Izquierdo LA, Smith JF, Blankstein J, Curet LB. Transvaginal sonographic measurement of cervical length: evaluation of twin pregnancies. *J Reprod Med* 1995;40:380-2.
32. Imseis HM, Albert TA, Iams JD. Identifying twin gestations at low risk for preterm birth with a transvaginal ultrasonographic cervical measurement at 24 to 26 weeks' gestation. *Am J Obstet Gynecol* 1997;177:1149-55.
33. Wennerholm UB, Holm B, Mattsby-Baltzer I, Nielsen T, Platz-Christensen J, Sundell G, Hosseini N, Hagberg H. Fetal fibronectin, endotoxin, bacterial vaginosis and cervical length as predictors of preterm birth and neonatal morbidity in twin pregnancies. *Br J Obstet Gynaecol* 1997;104:1398-404.
34. Crane JMG, Van den Hof M, Armson BA, Liston R. Transvaginal ultrasound in the prediction of preterm delivery: singleton and twin gestations. *Obstet Gynecol* 1997;90:357-63.
35. Goldenberg RL, Iams J, Miodovnik M, *et al.* The preterm prediction study: risk factors in twin gestations. *Am J Obstet Gynecol* 1996;175:1047-53.
36. The Collaborative Home Uterine Monitoring Study (CHUMS) Group. A multicenter randomized controlled trial of home uterine monitoring: active versus sham device. *Am J Obstet Gynecol* 1995;173:1120-7.
37. Dyson DC, Crites YM, Ray DA, Armstrong MA. Prevention of preterm birth in high-risk patients: the role of education and provider contact versus home uterine monitoring. *Am J Obstet Gynecol* 1991;164:756-62.
38. Rust OA, Perry KG, Andrew ME, Roberts WE, Martin RW, Morrison JC. Twins and preterm labor. *J Reprod Med* 1997;42:229-34.
39. Knuppel RA, Lake MF, Watson DL, Welch RA, Hill WC, Fleming AD, Martin RW, Bentley DL, Moenning RK, Morrison JC. Preventing preterm birth in twin gestation: home uterine activity monitoring and perinatal nursing support. *Obstet Gynecol* 1990;76:24S-27S.
40. Colton T, Kayne HL, Zhang Y, Heeren T. A meta-analysis of home uterine activity monitoring. *Am J Obstet Gynecol* 1995;173:1499-1505.
41. Lockwood CJ, Senyei A, Dische R, *et al.* Fetal fibronectin in cervical and vaginal secretions as a prediction of preterm delivery. *N Engl J Med* 1991;325:669-74.
42. Crane JP, Tomich PG, Kopta M. Ultrasonic growth patterns in normal and discordant twins. *Obstet Gynecol* 1980;55:678-83.
43. Erkkola R, Ala-Mello S, Piironen O, Abad L. Growth discordancy in twin pregnancies: a risk factor not detected by measurement of the biparietal diameter. *Obstet Gynecol* 1985;66:203-6.
44. Talbot GT, Goldstein RF, Nesbitt T, Johnson JL, Kay HH. Is size discordancy an indication for delivery of preterm twins? *Am J Obstet Gynecol* 1997;177:1050-4.
45. Chittkara U, Berkowitz GS, Levine R, *et al.* Twin pregnancy: routine use of ultrasound examinations in the prenatal diagnosis of intrauterine growth retardation and discordant growth. *Am J Perinatol* 1985;2:49-54.
46. Blickstein I, Friedman A, Caspi B, Lancet M. Ultrasonic prediction of growth discordancy by intertwin difference in abdominal circumference. *Int J Gynaecol Obstet* 1989;29:121-4.
47. Divon MY, Girz BA, Sklar A, Guidetti DA, Langer O. Discordant twins: a prospective study of the diagnostic value of real-time ultrasonography combined with umbilical artery velocimetry. *Am J Obstet Gynecol* 1989;161:757-60.
48. Hill LM, Guzik D, Cheney P, Boyles D, Nedzesky P. The sonographic assessment of twin growth discordancy. *Obstet Gynecol* 1994;84:501-4.
49. Storlazzi E, Vintzileos AM, Campbell WA, Nochimson DJ, Weinbaum PJ. Ultrasonic diagnosis of discordant fetal growth in twin gestation. *Obstet Gynecol* 1987;69:363-7.
50. Chamberlain P, Murphy M, Comerford FR. How accurate is antenatal sonographic identification of discordant birthweight in twins? *Eur J Obstet Gynecol Reprod Biol* 1991;40:91-6.
51. MacLean M *et al.* The ultrasonic assessment of discordant growth in twin pregnancies. *Ultrasound Obstet Gynecol* 1992;2:30-4.

52. Caravello JW, Chauhan SP, Morrison JC, Magann EF, Martin JN, Devoe LD. Sonographic examination does not predict twin growth discordance accurately. *Obstet Gynecol* 1997;89:529-34.
53. Cheung VYT, Bocking AD, Dasilva OP. Preterm discordant twins: What birth weight difference is significant? *Am J Obstet Gynecol* 1995;172:955-9.
54. Townsend RR, Simpson GF, Filly RA. Membrane thickness in ultrasound prediction of chorionicity of twin gestations. *J Ultrasound Med* 1988;7:3277-332.
55. D'Alton ME, Dudley DK. The ultrasonographic prediction of chorionicity in twin gestation. *Am J Obstet Gynecol* 1989;169:557-561.
56. Wood SL, Onge RS, Connors G, Elliot PD. Evaluation of the twin peak or lambda sign in determining chorionicity in multiple pregnancy. *Obstet Gynecol* 1996;88:6-9.
57. Spellcay WN, Handler A, Ferre CD. A case-control study of 1253 twin pregnancies from a 1982-1987 perinatal data base. *Obstet Gynecol* 1990;75:168-71.
58. Doyle PE, Beral V, Botting B, Wale CJ. Congenital malformations in twins in England and Wales. *J Epidemiol Community Health* 1990;45:43-8.
59. Edwards MS, Ellings JM, Newman RB, Menard MK. Predictive value of antepartum ultrasound examination for anomalies in twin gestations. *Ultrasound Obstet Gynecol* 1995;6:43-9.
60. Allen SR, Gray LS, Frentzen BH, Cruz AC. Ultrasonographic diagnosis of congenital anomalies in twins. *Am J Obstet Gynecol* 1991;165:1056-60.
61. Evans MI, Goldberg JD, Dommergues M *et al*. Efficacy of second-trimester selective termination for fetal abnormalities: International collaborative experience among the world's largest centers. *Am J Obstet Gynecol* 1994;171:90-4.
62. Ellings JM, Newman RB, Hulsey TC, Bivins HA, Keenan A. Reduction in very low birth weight deliveries and perinatal mortality in a specialized, multidisciplinary twin clinic. *Obstet Gynecol* 1993;81:387-91.
63. Dyson DC, Crites YM, Ray DA, Armstrong MA. Prevention of preterm birth in high risk patients: The role of education and provider contact versus home uterine monitoring. *Am J Obstet Gynecol* 1991;164:756-62.
64. Ewigman BG, Crane JP, Frigoletto FD, LeFebvre ML, Bain RP, McNellis D. Effect of prenatal ultrasound screening on perinatal outcome. *N Engl J Med* 1993;329:821-7.
65. LeFebvre ML, Bain RP, Ewigman BG, Frigoletto FD, Crane JP, McNellis D. A randomized trial of prenatal ultrasonographic screening: impact on maternal management and outcome. *Am J Obstet Gynecol* 1993;169:483-9.
66. Crane JP, LeFebvre ML, Winborn RC, *et al*. A randomized trial of prenatal ultrasonographic screening: impact on the detection, management and outcome of anomalous fetuses. *Am J Obstet Gynecol* 1994;171:392-9.
67. Saari-Kemppainen A, Karjalainen O, Ylösealo P, Heinonen OP. Ultrasound screening and perinatal mortality: controlled trial of systematic one-stage screening in pregnancy. *Lancet* 1990;336:387-91.
68. Waldenström U, Axelsson O, Nilsson S, *et al*. Effects of routine one-stage ultrasound screening in pregnancy: a randomized controlled trial. *Lancet* 1988;2:585-8.
69. Hughey MJ, Olive DL. Routine ultrasound scanning for the detection and management of twin pregnancies. *J Reprod Med* 1985;30:427-430.
70. Barrett JM, Staggs SM, Van Hoooydonk JE, Growdon JH, Killam AP, Boehm FH. The effect of type of delivery upon neonatal outcome in premature twins. *AJOG* 1982;143:360-7.
71. Chervenak FA. The controversy of mode of delivery in twins: the intrapartum management of twin gestation (part II). *Semin Perinatol* 1986;44-9.
72. Acker D, Lieberman M, Holbrook H, James O, Phillippe M, Edelin KC. Delivery of the second twin. *Obstet Gynecol* 1982;59:710-1.
73. Chervenak FA, Johnson RE, Berkowitz RL, Grannum P, Hobbins JC. Is routine cesarean section necessary for vertex-breech and vertex-transverse twin gestations? *AJOG* 1984;148:1-5.
74. Rabinovici J, Barkai G, Reichman B, Serr DM, Mashiach S. Randomized management of the second nonvertex twin: vaginal delivery or cesarean section. *AJOG* 1987;156:52-6.
75. Rabinovici J, Barkai G, Reichman B, Serr DM, Mashiach S. Internal podalic version with unruptured membranes for the second twin in transverse lie. *Obstet Gynecol* 1988;71:428-30.
76. Adam C, Allen AC, Baskett TF. Twin delivery: influence of the presentation and method of delivery on the second twin. *AJOG* 1991;165:23-7.
77. Fishman A, Grubb DK, Kovacs BW. Vaginal delivery of the nonvertex second twin. *AJOG* 1993;168:861-4.
78. Davison L, Easterling TR, Jackson JC, Benedetti TJ. Breech extraction of low-birth-weight second twins: can cesarean section be justified? *AJOG* 1992;166:497-502.
79. Gocke SE, Nageotte MP, Garite T, Towers CV, Dorchester W. Management of the nonvertex second twin: primary cesarean section, external version, or primary breech extraction. *AJOG* 1989;161:111-4.
80. Wells SR, Thorp JM, Bowes WZ. Management of the nonvertex second twin. *Surg Gynecol Obstet* 1991;172:383-5.
81. Chauhan SP, Roberts WE, McLaren RA, Roach H, Morrison JC, Martin JN. Delivery of the nonvertex second twin: breech extraction versus external cephalic version. *AJOG* 1995;173:1015-20.
82. Smith SJ, Zebrowitz J, Latta RA. Method of delivery of the nonvertex second twin: a community hospital experience. *J Matern-Fetal Med* 1997;6:146-50.
83. Levinsky E, Barrett JFR. Intrapartum management of twin gestations (Review). *Mat Fetal Med Rev*, 1999 (in press).
84. Adams DM, Chervenak FA. Intrapartum management of twin gestation. *Clin Obstet Gynecol* 1990;33:52-60.
85. Ismajovich B, Confino E, Sherzer A, Lidor A, David MP. Optimal delivery of nonvertex twins. *Mt Sinai J Med* 1985;52:106-109.
86. Poeschmann PP, Van Oppen CAC, Bruinse HW. Delayed interval delivery in multiple pregnancies: Report of three cases and review of the literature. *Obstet Gynecol Surv* 1992;47:139-47.
87. Rayburn WF, Lavin JP, Miodovnik M, Varner MW. Multiple gestation: Time interval between delivery of the first and second twins. *Obstet Gynecol* 1984;63:502-506.
88. Adam C, Allen AC, Baskett TF. Twin delivery: influence of presentation and method of delivery on the second twin. *AJOG* 1991;165:23-27.
89. Hecher K, Ville Y, Kypros H, Nicolaides KH. Color Doppler ultrasonography in the identification of communicating vessels in twin-twin transfusion syndrome and acardiac twins. *J Ultrasound Med* 1995;14:37-40.
90. Roberts, Mitchell JM. Fetal liver length in twin-twin transfusion syndrome. *Ultrasound Obstet Gynecol* 1997;9:30-4.
91. Sebire, D'Ercole C, Hughes K, Carvalho M, Nicolaides KH. Increased nuchal translucency thickness at 10-14 weeks of gestation as a predictor of severe twin-to-twin transfusion syndrome. *Ultrasound Obstet Gynecol* 1997;10:86-9.
92. Yamada A, Kasugai M, Ohno Y, Ishizuka T, Mizutani S, Tomoda Y. Antenatal diagnosis of twin-twin transfusion syndrome by Doppler ultrasound. *Obstet Gynecol* 1991;78:1058-61.
93. Saunders NJ, Snijders RJM, Nicolaides KH. Therapeutic amniocenteses in twin-twin transfusion syndrome appearing in the second trimester of pregnancy. *Am J Obstet Gynecol* 1992;166:820-4.
94. Nageotte MP, Hurwitz SR, Kaupke CJ, Vaziri ND, Pandian MR. Atriopeptin in the twin transfusion syndrome. *Obstet Gynecol* 1989;73:867-70.
95. Fries MH, Goldstein RB, Kilpatrick SJ, Golbus MS, Callen PW, Filly RA. The role of velamentous cord insertion in the etiology of twin-twin transfusion syndrome. *Obstet Gynecol* 1993;81:569-74.
96. Zosmer N, Bajoria R, Weiner E, Rigby M, Vaugham J, Fisk NM. Clinical and echocardiographic features of in-utero cardiac dysfunction in the recipient twin in twin-twin transfusion syndrome. *Dr Heart J* 1994; 72:74-9.
97. Naeye RL. Organ Abnormalities in a human parabiotic syndrome. *Am J pathol*.
98. Rausen AR, Seki M, Strauss L. Twin transfusion syndrome: review of 19 cases studied at one institution. *Journal of Pediatrics* 1995;66:613-28.
99. Sanders CH. The surgical pathologist examines the placenta. 235-88.
100. Sala MA, Matheus M. Placental characteristics in twin transfusion syndrome. *Arch Gynecol Obstet* 1989;246:51-6.
101. Blickstein I. The twin-twin transfusion syndrome. *Obstet Gynecol* 76(4):714-22.

102. Caglan MK, Kollee LAA. Determination of serum ferritin in the evaluation of iron depletion and iron over load in chronic twin-to-twin transfusion syndrome. *J Perinat Med* 1989;17:357-9.
103. Wittman BK, Baldwin VJ, Nichol B. Antenatal diagnosis of twin transfusion syndrome by ultrasound. *Obstet Gynecol* 1981;58:123-6.
104. Danskin FH, Neilson JP. Twin-to-twin transfusion syndrome: what are appropriate diagnostic criteria? *Am J Obstet Gynecol* 1989;161:365-9.
105. Fisk NM, Borrell A, Hubinont C, Tannirandorn Y, Nicolini U, Rodeck CH. Fetofetal transfusion syndrome: do the neonatal criteria apply in utero? *Arch Disease Child* 1990;65:657-61.
106. Brennan JN, Diwan RV, Rosen MG, Bellon EM. Fetofetal transfusion syndrome: prenatal ultrasonographic diagnosis. *Radiology* 1982;143:535-6.
107. Brown DL, Genson CB, Driscoll SG, Doubilet PM. Twin-twin transfusion syndrome: sonographic findings. *Radiology* 1989;170:61-3.
108. Chescheir, NC, Seeds JW. Polyhydramnios and oligohydramnios in twin gestation. *Obstet Gynecol* 1988;71:882-4.
109. McCulloch K. Neonatal problems in twins. *Clinics Perinat* 1988; 15(1):141-58.
110. Nicosia RF, Krouse TB, Mobini J. Congenital aortic intimal thickening: its occurrence in a case of twin-transfusion syndrome. *Arch Pathol Lab Med* 1981;105:247-9.
111. Alteminia AM, Vassallo J, Billis A. Congenital focal glomerular lesions in only one monozygous twins related to probable twin transfusion syndrome. *Histopathology* 1986;10:991-4.
112. Mahony BS, Petty CN, Nyberg DA, Luthy DA, Hickok DE, Hirsch HJ. The "stuck twin" phenomenon: ultrasonographic findings, pregnancy outcome, and management with serial amniocenteses. In prenatal detection of congenital toxoplasmosis, from *Am J. Obstet Gynecol* 1990;163:1513-22.
113. Michael G, Pinette, MD, Yuqun Pan, MM, Sheila Gerry Pinette, RPAC, and Phillip G. Stubblefield, MD, Treatment of Twin-Twin Transfusion Syndrome, from *Obstet Gynecol* 1993;82:841-6.
114. Elliott JP, Urig MA, Clewell WH. Aggressive therapeutic amniocentesis for treatment of twin-twin transfusion syndrome. *Obstet Gynecol* 1991;77:537-40.
115. Saunders, Snijders RJM, Nicolaidis KH. Therapeutic amniocenteses in twin-twin transfusion syndrome appearing in the second trimester of pregnancy. *Am J Obstet Gynecol* 1992;166:820-4.
116. Lopriore E, Vandenbussche FPHA, Tiersma ESM, de Beaufort AJ, de Leeuw JP. Twin-to-twin transfusion syndrome: new perspectives. *J Pediatr* 1995;127(5):675-80.
117. Elliott JP, Sawyer AT, Radin TG, Strong RE. Large-volume therapeutic amniocentesis in the treatment of hydramnios. *Obstet Gynecol* 1994;84:1025-7.
118. Bebbington MW, Wilson RD, Machan L, Wittmann BK. Selective fetocide in twin transfusion syndrome using ultrasound-guided insertion of thrombogenic coils. *Fetal Diagn Ther* 1995;10(1):32-6.
119. Dommergues M, Mandelbrot L, Delezoide AI, et al. Twin-to-twin transfusion syndrome: selective fetocide by embolization of the hydropic fetus. *Fetal Diagn Ther* 1995;10:26-31.
120. Mari G. Amnioreduction in twin-twin transfusion syndrome: a multi-center registry, evaluation of 579 procedures. *Am J Obstet Gynecol* 1998;178:S28.
121. Challis D, Dockrill K, Barrett J, et al. Aggressive therapeutic amniocentesis in the management of twin-twin transfusion syndrome. *Am J Obstet Gynecol* 1997;176:S85.
122. Saade G, Belfort M, Berry D, et al. Amniotic septostomy for the treatment of twin oligohydramnios-polyhydramnios sequence. *Fetal Diagn Ther* 1998;13:86-93.
123. Ville Y, Hyett J, Hecher K, Nicolaidis K. Preliminary experience with endoscopic laser surgery for severe twin-twin transfusion syndrome. *N Engl J Med* 1995;332:224-7.
124. Delia JE, Kuhlmann RS, Harstad TW, Cruikshank DP. Fetoscopic laser ablation of placental vessels in severe previable twin-twin transfusion syndrome. *Am J Obstet Gynecol* 1995;172:1202-11.
125. Ville Y, Hecher K, Gagnon A, Sebire N, Hyett J, Nicolaidis K. Endoscopic laser coagulation in the management of severe twin-to-twin transfusion syndrome. *Br J Obstet Gynaecol* 1998;105:446-53.
126. Nicolaidis K, Pettersen H. Fetal therapy. *Obstet Gynecol* 1994;6:468-471.
127. Radesat A, Thomassen PA. Acute polyhydramnios in twin pregnancy: a retrospective study with special reference to therapeutic amniocentesis. *Acta Obstet Gynecol Scand* 1990;69:297-300.
128. Wittmann BK, Farquharson DF, Thomas WDS, Baldwin VJ, Wadsworth LD, The role of feticide in the management of severe twin transfusion syndrome. *Am J Obstet. Gynecol* 1986;155:1023-6.
129. Machin GA, Keith LG. Can twin-to-twin transfusion syndrome be explained, and how is it treated? *1998;41(1):105-15.*
130. Feingold M, Cetrulo CL, Newton ER, Weiss J, Shaker C, Shmoys S. Serial amniocenteses in the treatment of twin to twin transfusion complicated with acute polyhydramnios. *ACTA Genet Med. Gemellol* 1986;35:107-13.
131. Mahony BS, Petty CN, Nyberg DA, Luthy DA, Hickok DE, Hirsch JH. The "stuck twin" phenomenon: ultrasonographic findings, pregnancy outcome, and management with serial amniocenteses. *Am J Obstet Gynecol* 1990;163:1513-22.
132. Elliott JP, Urig MA, Clewell WH. Aggressive therapeutic amniocentesis for treatment of twin-twin transfusion syndrome. *Obstet Gynecol* 1991;77(4):537-40.
133. Urig MA, Sijmpson GF, Elliott JP, Clewell WH. Twin-twin transfusion syndrome: the surgical removal of one twin as a treatment option. *Fetal Therapy* 1988;3:185-8.
134. De Lia JE, Cruikshank DP, Keye WR. Fetoscopic neodymium:Yag laser occlusion of placental vessels in severe twin-twin transfusion syndrome. *Obstet Gynecol* 1990;75(6):1046-53.
135. Danziger RW, Chir B. Twin pregnancy with acute hydramnios treated by paracentesis uteri. *Br Med J* 1948;2:205-6.
136. Berry D, Montgomery L, Johnson A, Saade G, Moise K. Amniotic septostomy for the treatment of the stuck twin sequence. *Am J Obstet Gynecol* 1997;176:519 (abstr).
137. Porreco R, Barton S, Haverkamp A. Occlusion of umbilical artery in acardiac, acephalic twin. *Lancet* 1991;337:326-7.
138. Golan A, Amit A, Baram, David MP. Unusual cord intertwining in monoamniotic twins. *J Obstet Gynaec* 1982;22:165-7.
139. Salerno LJ. Monoamniotic twinning: a survey of the American literature since 1935 with a report of four new cases. *Obstet Gynecol* 1959;14:205-13.
140. Georgeson S, Sonnenberg FA, Feingold M, Pauker SG. Twisted sisters: when is the optimal time for delivery? *Med Decision Making* 1990;10:295-301.
141. Dorum A, Nesheim BI. Monochorionic monoamniotic twins: the most precarious of twin pregnancies. *Acta Obstet Gynecol Scand*, 1991;70(4-5):381-3.
142. Finberg NJ, Clewell WH. Definitive prenatal diagnosis of monoamniotic twins. Swallowed amniotic contrast agent detected in both twins on sonographically selected CT images. *J Ultrasound Med* 1991;10(9):513-6.
143. Sargent SK, Young W, Crow P, Simpson W. CT amniography: value in detecting a monoamniotic pair in a triplet pregnancy. *AJR* 1991;156(3): 559-60.
144. Tessen JA, Zlatnik FJ. Monoamniotic twins: a retrospective controlled study. *Obstet Gynecol* 1991;77(6):832-4.
145. Carr SR, Aronson MP, Coustan DR. Survival rates of monoamniotic twins do not decrease after 30 weeks' gestation. *Amer J Obstet Gynecol* 1990;163(3):719-22.
146. Rodis JF, Vintziliou AM, Campbell WA, Deaton JL, Funia F, Nochimson DJ. Antenatal diagnosis and management of monoamniotic twins. *Amer J Obstet Gynecol* 1987;157(5):1255-7.
147. Griffith HB. Monoamniotic twin pregnancy. *Brit J Clin Pract* 1986; 40(7):294-7.
148. Kassam SH, Tompkins MG. Monoamniotic twin pregnancy and modern obstetrics: report of a case with a peculiar cord complication. *Diagnost Gynecol Obstet* 1980;2(3):213-20.
149. Langer B, Boudier E, Gasser B, Christmann D, Messer J, Schlaeder G. Antenatal diagnosis of brain damage in the survivor after the second trimester death of a monochorionic monoamniotic co-twin: case report and literature review. *Fetal Diagn Ther* 1997;12(5): 286-91.
150. Rodis JF, McIlveen PF, Egan JF, Borgida AF, Turner GW, Campbell WA.

- 
- Monoamniotic twins: improved perinatal survival with accurate prenatal diagnoses and antenatal fetal surveillance. *Amer J Obstet Gynecol* 1997;177(5):1046-9.
151. Peek MJ, McCarthy A, Kyle P, Sepulveda W, Fisk NM. Medical amnioreduction with sulindac to reduce cord complications in monoamniotic twins. *Amer J Obstet Gynecol*, 1997;176(2):334-6.
152. Ritossa M, O'Loughlin J. Monoamniotic twin pregnancy and cord entanglement: a clinical dilemma (review, 12 refs).
153. Dubecq F, Dufour P, Vinatier D, *et al.* Monoamniotic twin pregnancies: review of the literature and a case report with vaginal delivery. *Eur J Obstet Gynecol Reprod Biol* 1996;66(2):183-6.
154. Benirschke K. The biology of the twinning process: how placentation influences outcome. *Sem Perinatol* 1995;19(5):342-50.
155. Strohhahn K, Dattel BJ. Pitfalls in the diagnosis of nonconjoined monoamniotic twins. *J Perinatol* 1995;15(6):484-93.
156. Aisenbrey GA, Catanzarite VA, Hurley TJ, Spiegel JH, Schrimmer DB, Mendoza A. Monoamniotic and pseudomonoamniotic twins: sonographic diagnosis, detection of cord entanglement, and obstetric management. *Obstet Gynecol* 1995;86(2):218-22.