

Percutaneous endoscopic laser thoracic discectomy

1 Guidance

- 1.1 Current evidence on the safety and efficacy of percutaneous endoscopic laser thoracic discectomy does not appear adequate for this procedure to be used without special arrangements for consent and for audit or research.
- 1.2 Clinicians wishing to undertake percutaneous endoscopic laser thoracic discectomy should take the following action.
 - Inform the clinical governance leads in their Trusts.
 - Ensure that patients understand the uncertainty about the procedure's safety and efficacy and provide them with clear written information. Use of the Institute's *Information for the Public* is recommended.
 - Audit and review clinical outcomes of all patients having percutaneous endoscopic laser thoracic discectomy.
- 1.3 Further research will be useful in reducing the current uncertainty and clinicians are encouraged to collect longer-term follow-up data. The Institute may review the procedure upon publication of further evidence.

2 The procedure

2.1 Indications

- 2.1.1 Percutaneous endoscopic laser thoracic discectomy is used to treat symptomatic thoracic disc herniation. This occurs when a portion of the intervertebral disc protrudes into the spinal canal and impinges on a nerve root. Symptoms include back pain, radicular pain, nondermatomal leg pain, bladder dysfunction and lower extremity weakness. If left untreated, serious neurological sequelae may occur.
- 2.1.2 Standard discectomy for thoracic disc herniation may be either by open posterolateral or anterior approaches. A percutaneous endoscopic approach may lessen the morbidity associated with the procedure by allowing access and visualisation of the anterior and lateral aspects of the disc. The choice of approach will depend upon the characteristics of the disc herniation and the surgeon's experience with the above techniques.

2.2 Outline of the procedure

- 2.2.1 Percutaneous endoscopic laser thoracic discectomy is usually done under local anaesthesia through a small incision in the back, using X-ray monitoring. A needle is introduced into the centre of the affected intervertebral disc. A guidewire is passed through the needle, followed by small instruments, which are used to remove some disc material. A Holmium-YAG laser is then

Interventional Procedure Guidance 61

This guidance is written in the following context:

This guidance represents the view of the Institute which was arrived at after careful consideration of the available evidence. Health professionals are expected to take it fully into account when exercising their clinical judgement. This guidance does not, however, override the individual responsibility of health professionals to make appropriate decisions in the circumstances of the individual patient, in consultation with the patient and/or guardian or carer.

introduced and laser energy is used to destroy more of the disc. Debris is removed by surgical instruments. The patient's neurological status is monitored throughout.

2.3 Efficacy

- 2.3.1 No controlled studies were identified. The studies identified provided little detail of study design and outcomes. In one study 96% (96/100) of patients reported 'good-to-excellent results/symptomatic relief', but the meaning of this was not defined. The average time to return to work in this study was 10 days. For more details, refer to the sources of evidence (see below).
- 2.3.2 One Specialist Advisor commented that there was no evidence to support the efficacy of the procedure, and that the procedure was difficult to master.

2.4 Safety

- 2.4.1 No operative or postoperative complications were reported in the studies identified. However, these studies provided little detail of study design and outcomes.
- 2.4.2 One Specialist Advisor considered that this procedure had the potential for serious neurological complications, and was concerned about risks to patients while surgeons learnt the procedure. This Advisor also thought that the procedure could result in nerve injury.

2.5 Other comments

- 2.5.1 This decision relates to the procedure when used in isolation (for example, to treat degenerative disc disease). No judgement is made regarding the use of this procedure as part of a larger operation, such as the treatment of scoliosis.
- 2.5.2 Appropriate patient selection for this procedure is important and may be difficult.

Andrew Dillon
Chief Executive
May 2004

Information for the Public

The Institute has produced information describing its guidance on this procedure for patients, carers and those with a wider interest in healthcare. It explains the nature of the procedure and the decision made, and has been written with patient consent in mind. This information is available, in English and Welsh, from www.nice.org.uk/IPG061publicinfo.

Sources of evidence

The evidence considered by the Interventional Procedures Advisory Committee is described in the following document.

Interventional procedure overview of percutaneous endoscopic laser thoracic discectomy, October 2002.

Available from: www.nice.org.uk/ip176overview

Ordering information

Copies of this guidance can be obtained from the NHS Response Line by telephoning 0870 1555 455 and quoting reference number N0580. *Information for the Public* can be obtained by quoting reference number N0581 for the English version and N0582 for a version in English and Welsh.

The distribution list for this guidance is available on the NICE website at URL www.nice.org.uk/IPG061distributionlist

Published by the National Institute for Clinical Excellence, May 2004 ISBN: 1-84257-644-5

© National Institute for Clinical Excellence May 2004. All rights reserved. This material may be freely reproduced for educational and not for profit purposes within the NHS. No reproduction by or for commercial organisations is permitted without the express written permission of the Institute.

National Institute for Clinical Excellence

MidCity Place, 71 High Holborn, London WC1V 6NA, website: www.nice.org.uk

N0580 1P 20k May 04 (HOBBS)